

Understanding the Underlying Event

G.G. Barnaföldi in collaboration with A.N. Mishra, G. Paic, and G. Bíró

Support: *Hungarian OTKA grants, NK123815, K135515 Wigner GPU Laboratory*
Ref: *arXiv:2108.13938*

ELTE TTK ElmFiz Seminar, Budapest, 19th October 2021



ELKH | Eötvös Loránd
Research Network

Outline

1) Earlier studies

- What is UE? Why is this important for in HEP?
→ theory, experiment, measures

2) New developments on UE

- Angular properties measures
→ multiplicity, p_T spectra, parameter derivatives
→ Tsallis thermometer


3) Comparison to event shape variable

- Sphericity measures and cross check


→ **Conclusions: Extended UE definition**




Anatomy of a proton-proton event




Anatomy of a proton-proton event



Anatomy of a proton-proton event




Anatomy of a proton-proton event



So what Underlying Event is?

- **Theoretical point:**
 - Mainly non-perturbative QCD effect
 - Initial & final state radiation
 - Multiple parton interaction
 - Color Reconnection (CR)
 - intrinsic k_T
 - Hadronization





So what Underlying Event is?

- **Theoretical point:**

- Mainly non-perturbative QCD effect
 - Initial & final state radiation
 - Multiple parton interaction
 - Color Reconnection (CR)
 - intrinsic k_T
 - Hadronization


- **Experimental point**

- Pedestal-like effects
 - Activity in the event over MB
 - Beam remnants (pile up)
 - Trigger bias (jet criterion)





Earlier studies, motivation

Geometrical structure of an event



Geometrical structure of an event




How to separate jet & UE?

- **Jet finding & elimination:**

- Surrounding Band (SB method), Find a jet, THEN define SBs
- IF SB_1 and SB_2 are equal, THEN eliminate the jet
 - expensive (high statistics)
 - sensitive to cuts

- **Correlation & background**

- Traditional method by CDF
 - brute force
 - geometry info only




How to separate jet & UE?

- **Jet finding & elimination:**

- Surrounding Band (SB method), Find a jet, THEN define SBs
- IF SB_1 and SB_2 are equal, THEN eliminate the jet
 - expensive (high statistics)
 - sensitive to cuts

- **Correlation & background**

- Traditional method by CDF
 - burte force
 - geometry info only



How to quantify & compare events?

- **Transverse sphericity:**


$$S_0 = \frac{\pi^2}{4} \left(\frac{\sum_i |\vec{p}_{T,i} \times \hat{n}|}{\sum_i p_{T,i}} \right)^2$$

- **Thrust:**

$$T_{\min} \equiv \frac{\sum_i |\vec{p}_{T,i} \cdot \hat{n}_m|}{\sum_i p_{T,i}}$$

→ **NO** need for jet fin

→ **Momentum & geometry** infos



How to quantify & compare events?

- **Precise spectra description**

- from low- to high- p_T

$$f(m_T) = A \cdot \left[1 + \frac{q-1}{T_s} (m_T - m) \right]^{-\frac{1}{q-1}}$$

- in multiplicity classes (pp, pA, AA)


$$\left. \frac{dN_{\text{ch}}}{dy} \right|_{u=0} = 2\pi A T_s \left[\frac{(2-q)m^2 + 2mT_s + 2T_s^2}{(2-q)(3-2q)} \right] \times \left[1 + \frac{q-1}{T_s} m \right]^{-\frac{1}{q-1}}$$

- **With PID:**

$$\pi^\pm, K^\pm, K_s^0, K^{*0}, p(\bar{p}), \Phi, \Lambda, \Xi^\pm, \Sigma^\pm, \Xi^0, \Omega$$

- **Wide range:**

	pp	pA	AA
CM energy (GeV)	7000, 13000	5020	130-5020
Multiplicity range	2.2-25.7	4.3-45	13.4-2047



How to quantify & compare events?

- **QCD-inherited scaling properties**




$$f(m_T) = A \cdot \left[1 + \frac{q-1}{T_s} (m_T - m) \right]^{-\frac{1}{q-1}}$$

- Parameter scaling with \sqrt{s} & multiplicity

$$A(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = A_0 + A_1 \ln \frac{\sqrt{s_{NN}}}{m} + A_2 \langle N_{ch}/\eta \rangle$$

$$T(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = T_0 + T_1 \ln \frac{\sqrt{s_{NN}}}{m} + T_2 \ln \ln \langle N_{ch}/\eta \rangle,$$

$$q(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = q_0 + q_1 \ln \frac{\sqrt{s_{NN}}}{m} + q_2 \ln \ln \langle N_{ch}/\eta \rangle,$$



How to quantify & compare events?

- QCD-inherited scaling properties**

$$f(m_T) = A \cdot \left[1 + \frac{q-1}{T_s} (m_T - m) \right]^{-\frac{1}{q-1}}$$

- Parameter scaling with \sqrt{s} & multiplicity

$$A(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = A_0 + A_1 \ln \frac{\sqrt{s_{NN}}}{m} + A_2 \langle N_{ch}/\eta \rangle$$


$$T(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = T_0 + T_1 \ln \frac{\sqrt{s_{NN}}}{m} + T_2 \ln \ln \langle N_{ch}/\eta \rangle$$

$$q(\sqrt{s_{NN}}, \langle N_{ch}/\eta \rangle, m) = q_0 + q_1 \ln \frac{\sqrt{s_{NN}}}{m} + q_2 \ln \ln \langle N_{ch}/\eta \rangle$$

- Thermodynamical consistency**


$$P = g \int \frac{d^3 p}{(2\pi)^3} T f, \quad N = nV = gV \int \frac{d^3 p}{(2\pi)^3} f^q,$$

$$s = g \int \frac{d^3 p}{(2\pi)^3} \left[\frac{E-\mu}{T} f^q + f \right], \quad \varepsilon = g \int \frac{d^3 p}{(2\pi)^3} E f$$




New development to understand UE

Angular structure of an event




Standard CDF definition

Angular structure of an event




Standard CDF definition




Case I: Opening angle


Angular structure of an event



Standard CDF definition



Case I: Opening angle




Case II: Sliding angle

Angular structure of an event

- **Case I: Opening angle**


- We open $\Delta\phi$ angle in steps of 20° . The binning starts from -10° to 10° and the last bin covers full azimuthal space i.e. -180° to 180° (MB). Case I is useful to investigate the evolution of the thermodynamical observables of the system.



Case I: Opening angle

- **Case II: sliding angle**


- We make slices of the $\Delta\phi$ of size 20° . In this case, the results for the first bin 0 to 20° are reported in two ways: including and excluding the leading particle in the result. Case II is a tool for exploring the geometrical structure of the Underlying Event.




Case II : Sliding angle

The simulated data

- **PYTHIA_v8240 Monash 2013 tune**
 - 1 billion non-diffractive collisions of pp
 - C.m. energy: $\sqrt{s} = 13$ TeV
 - Includes $2 \rightarrow 2$ hard scattering process, followed by initial and final state parton showering, multiparton interactions, and the final hadronization process.
 - The events having at least three primary charged particle with transverse
 - Min. momentum: $p_T > 0.15$ GeV/c
 - Pseudorapidity: $|\eta| < 0.8$
 - UE: Color Reconnection (CR, Multiple Parton Interaction (MPI)




Case I: Opening angle "Pacman"



Case I: Opening angle

Case I: Opening angle "Pacman"




Case I: Opening angle


Case I: Multiplicity/MB

- **PYTHIA multiplicity with opening angle**

- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations
- More multiplicity in the NS
- Getting flat in NS+TS
- NS+TS+AS are mainly flat as reaching MB




Case I: Opening angle



Case I: Multiplicity/MB

- **PYTHIA multiplicity with opening angle**


- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations
- More multiplicity in the NS
- Getting flat in NS+TS
- NS+TS+AS are mainly flat as reaching MB




Case I: p_T spectrum

- **PYTHIA spectra with opening angle**

- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations
- Low p_T varies (T)
- High p_T is constant (q)
- Full opening is MB




Case I: Opening angle




Case I: p_T spectrum

- **PYTHIA spectra with opening angle**

- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations
- Low p_T varies (T)
- High p_T is constant (q)
- Full opening is MB




Case I: Opening angle







Case I: Tsallis fit parameters

- PYTHIA spectra with opening angle**

- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations (red line)
- Opening the angle \rightarrow constant T , lowest at MB
- Opening the angle \rightarrow constant q
- Multiplicity $\sim A$
- Full opening is MB



Case I: Opening angle




Case I: derivatives of the parameters

- PYTHIA spectra parameter derivatives with opening angle**





- PYTHIAs model UE: CR & MPI
- Opening the angle \rightarrow constant T & q

$$\frac{\delta T_s}{\delta(\Delta\phi)} \rightarrow 0 \quad \& \quad \frac{\delta q}{\delta(\Delta\phi)} \rightarrow 0$$

- No change beyond NS
- Multiplicity $\sim A$
- Full opening is MB



Case I: Opening angle




Case II: Sliding angle "cake slices"



Case II : Sliding angle

Case II: Sliding angle "cake slices"




Case II : Sliding angle


Case II: Multiplicity/MB

- **PYTHIA multiplicity with sliding angle**

- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations
- More multiplicity as NS
- TS & AS are mainly flat
- With leading particle deviation is increased




Case II : Sliding angle



Case II: Multiplicity/MB

- **PYTHIA multiplicity with sliding angle**


- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations
- More multiplicity as NS
- TS & AS are mainly flat
- With leading particle deviation is increased




Case II: p_T spectrum

- **PYTHIA spectra with sliding angle**

- PYTHIA model UE: CR & MPI
- Good fits with the parametrizations
- Low p_T is constant (T)
- High p_T varies (q)
- NS/AS are similar
- Need to consider w/o leading particle




Case II : Sliding angle




Case II: p_T spectrum

- **PYTHIA spectra with sliding angle**

- PYTHIA model UE: CR & MPI
- Good fits with the parametrizations
- Low p_T is constant (T)
- High p_T varies (q)
- NS/AS are similar
- Need to consider w/o leading particle




Case II : Sliding angle







Case II: Tsallis fit parameters

- PYTHIA spectra with sliding angle**

- PYTHIA's model UE: CR & MPI
- Good fits with the parametrizations (red line)
- NS \rightarrow highest T
- NS/AS \rightarrow highest q
- TS \rightarrow constant q, T
- Multiplicity $\sim A$



Case II : Sliding angle



Case II: derivatives of the parameters


- PYTHIA spectra parameter derivatives with sliding angle**

- PYTHIA's model UE: CR & MPI
- TS (+AS) → constant T & q





$$\frac{\delta T_s}{\delta(\Delta\phi)} \neq 0 \quad \& \quad \frac{\delta q}{\delta(\Delta\phi)} \neq 0 \quad (\text{for NS \& AS})$$

$$\frac{\delta T_s}{\delta(\Delta\phi)} \approx 0 \quad \& \quad \frac{\delta q}{\delta(\Delta\phi)} \approx 0 \quad (\text{for TS})$$

- NS → highest T
- NS/AS → highest q
- Multiplicity ~ A




Case II : Sliding angle



On the Tsallis-thermometer

- **Case I: opening angle**

- Need UE in PYTHIA → CR & MPI
- NS → highest T, lowest q
- TS/AS → constant q, lowering T
- MB → constant q, lowest T




On the Tsallis-thermometer

- **Case I: opening angle**

- Need UE in PYTHIA → CR & MPI
- NS → highest T, lowest q
- TS/AS → constant q, lowering T
- MB → constant q, lowest T

- **Case II: sliding angle**

- Need UE in PYTHIA → CR & MPI
- NS (with leading) is fully different highest T & highest q
- Beyond NS T is getting constant → Wider range of UE, than in CDF





Cross-check with event shape variable

Event shape variable: sphericity

Simple 2-component model


- Isotrope: flat low p_T distribution
- Jet: flat high p_T distribution



Event shape variable: sphericity

Simple 2-component model

- Isotrope: flat low p_T distribution
- Jet: flat high p_T distribution





Event shape variable: sphericity

Simple 2-component model

- Isotrope: flat low p_T distribution
- Jet: flat high p_T distribution

Sphericity definition



$$S_0 = \frac{\pi^2}{4} \left(\frac{\sum_i |\vec{p}_{T_i} \times \hat{n}|}{\sum_i p_{T_i}} \right)^2$$



→ Event selection based on sphericity classes is available in ALICE

Case II: Spherocity vs. Tsallis thermometer



- Spherocity relative to the MB defines wider UE



→ Wider range of UE [40,140], than in CDF [60,120]

Case II: Spherocity vs. Tsallis thermometer


- Spherocity relative to the MB defines wider UE
- Tsallis-thermometer presents the same



→ Wider range of UE [40,140], than in CDF [60,120]

Case II: Spherocity vs. Tsallis thermometer



- Spherocity relative to the MB defines wider UE
- Tsallis-thermometer presents the same



→ Wider range of UE [40,140], than in CDF [60,120]

Case II: Parameters in spherocity classes

- **PYTHIA spectra with sliding angle in S_0 classes**
 - The more jetty the event, the angular variation is stronger.
 - Minimal activity (lowest q & T values are in the isotropic case.



→ Isotropic event are closer to UE, activity is more than MB

Conclusions

- **Could we understand UE?**
 - Not yet, but getting closer by quantifying them
 - Model UE: PYTHIA (CR, MPI), HIJING (minijet)
 - UE properties has been charaterized
 - Tsallis-Pareto fits well in narrow slices
- **To take away...**
 - Tsallis-thermometer present wider UE
 - In degrees CDF: [60,120] → [40,140]
 - Event shape classification support the model




Conclusions

- **Could we understand UE?**
 - Not yet, but getting closer by quantifying them
 - Model UE: PYTHIA (CR, MPI), HIJING (minijet)
 - UE properties has been charaterized
 - Tsallis-Pareto fits well in narrow slices
 - **To take away...**
 - Tsallis-thermometer present wider UE
 - In degrees CDF: [60,120] → [40,140]
 - Event shape classification support the model
- **Next stage can be in a more complex system**




BACKUP

Relative spherocity/MB Case I and II




(a) Case I




(b) Case II

Case I: Parameters vs spherocity



Spherocity model with multiplicity



Thermodynamical consistency?



Thermodynamical consistency: fulfilled up to a high degree

$$P = g \int \frac{d^3p}{(2\pi)^3} T f,$$

$$N = nV = gV \int \frac{d^3p}{(2\pi)^3} f q,$$

$$s = g \int \frac{d^3p}{(2\pi)^3} \left[\frac{E - \mu}{T} f q + f \right],$$

$$\varepsilon = g \int \frac{d^3p}{(2\pi)^3} E f$$



Compare EoS to data: Lattice QCD (parton) & Biró-Jakovác parton-hadron

