



A Deep Learning Based Estimator for Elliptic Flow in Heavy ion Collisions

N. Mallick^{1,†}, S. Prasad¹, A. N. Mishra², R. Sahoo^{1,3}, and G. G. Barnaföldi²

¹Department of Physics, Indian Institute of Technology Indore, Simrol, Indore 453552, India,


²Wigner Research Center for Physics, 29-33 Konkoly-Thege Miklós Str, H-1121 Budapest, Hungary, and

³CERN, CH 1211, Geneva 23, Switzerland



1. Introduction

- Transverse collective flow is an important observable in studying Quark-Gluon Plasma (QGP) [1]
- Heavy-ion collisions observe significant elliptic flow (v_2) [2]
- Estimation of reaction plane angle (ψ_R) is non-trivial [3]
- Deep Neural Networks are well suited for mapping complex nonlinear functions [4]



$$v_n = \langle \cos[n(\phi - \psi_R)] \rangle$$

First Deep Learning based estimator for elliptic flow [5]

Input Vector
(3072)

Input Layer
(128, ReLU)

Hidden Layer (I)
(256, ReLU)


Hidden Layer (II)
(256, ReLU)

Hidden Layer (III)
(256, ReLU)


Output Vector
(1, Linear)

2. Deep Learning Estimator


- $(\eta - \phi)$ space as the primary input space
- p_T , mass, and an energy term as the secondary inputs
- 32 × 32 bins each, three such input layers
- Training with Pb-Pb collisions, $\sqrt{s_{NN}} = 5.02$ minimum bias events, simulated with a multiphase transport model (AMPT)
- Early stopping callback to ensure minimal overfitting
- Mean Absolute Error ($\Delta v_2 = 0.0073$) on 10K testing data
- For i^{th} event, and j^{th} feature, the feature value $F_{i,j} \leftarrow F_{i,j} + X_{i,j}/w$, where $X_{i,j} \in (-\sigma_j, \sigma_j)$.
- σ_j = standard deviation, w = noise parameter




Centrality Dependence



3. Results



- From minimum bias training to estimation of centrality and p_T dependence of v_2
- Comparison of prediction with simulation and experimental results
- Estimation of v_2 for different collision systems



4. Summary

- Final state particle kinematic information as input
- Estimator preserves the centrality, and p_T dependence of v_2
- Excellent prediction accuracy against noisy simulation
- Applicable to both RHIC and LHC energy
- v_2 for identified particles, and n_q scaling under study

References

- U. Heinz and R. Snellings, Ann. Rev. Nucl. Part. Sci. **63**, 123 (2013).
- J. Adams et al. [STAR], Phys. Rev. Lett. **92**, 052302 (2004).
- A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C **58**, 1671 (1998).
- K. Hornik, M. Stinchcombe and H. White, Neural Netw. **2**, 359 (1989).
- N. Mallick, S. Prasad, A. N. Mishra, R. Sahoo, and G. G. Barnaföldi, [arXiv:2203.01246 [hep-ph]], and references therein.

E-mail: Neelkamal.Mallick@cern.ch